arXiv:0908.1074v1 [math.PR] 7 Aug 2009

FUNCTIONAL LIMIT THEOREMS FOR LEVY PROCESSES AND THEIR ALMOST SURE VERSIONS

By ELENA PERMIAKOVA (Kazan)
Chebotarev Inst. of Mathematics and Mechanics, Kazan State University
Universitetskaya 17, 420008 Kazan
e-mail: epermiakova@mail.ru

Abstract. In the paper it proved a criterion of convergences in distribution in Skorohod space. This criterion is applied to some special Levy processes. The almost sure version of limit theorems for this processes are obtained.

2000 AMS Mathematics Subject Classification. 60F05 Central limit and other weak theorems, 60F15 Strong theorems.

Key words and phrases: functional limit theorem, almost sure limit theorem, α-stable random process, Levy random process.

1 Introduction.

Consider the sequence of the random processes

$$X_n(x) = \frac{V(s_n x)}{a_n} - x b_n, \quad x \in [0,1], \quad n \in \mathbb{N} \tag{1}$$

where $s_n \to \infty$ as $n \to \infty$, $V = V(x)$, $x \in [0,\infty)$ is a Levy process and a_n, $b_n \in \mathbb{R}$. In this paper we solve the question of approximation α-stable processes by Levy processes of type (1).

In Section 2 we prove the criterion of convergence of sequence Levy processes V_n in distribution in the Skorohod space $D[0,1]$ (Theorem 1). As a corollary we obtain a theorem of the convergence in distribution of compound Poisson processes. This corollary can be used from approximation the Levy process by random sums with the Poisson index summation.

In Section 3 we introduce conditions for the process V under which the sequence X_n converges in distribution in $D[0,1]$ to a α-stable Levy process Y_α, $0 < \alpha < 2$ (Theorem 2) and conditions for process V under which X_n converges in distribution in $D[0,1]$ to a Wiener random process W (Theorem 4). In particular we consider $a_n = n^{1/\alpha}$ and specify conditions for the process V under which X_n converges in distribution to Y_α, $0 < \alpha < 2$ (Theorem 3).

In Section 4 it proved almost sure versions of the limit theorems from Section 3. We describe the sequences (s_n) such that the sequence of measures

$$Q_n(\omega) = \frac{1}{\ln n} \sum_{k=1}^{n} \frac{1}{k} \delta_{X_k(t)(\omega)}, \quad n \in \mathbb{N} \tag{2}$$

where δ_x denotes the measure of unit mass, concentrated in the point x, converges weakly to a distribution of Y_α or W in $D[0,1]$ for almost all $\omega \in \Omega$.

The proofs of our results are based on the criterion for integral type almost sure version of a limit theorem which was obtained in Chuprunov and Fazekas [4].

2 Functional limit theorems.

We will denote by $\xrightarrow{d}$ the convergence in distribution, by $\overset{d}{=}$ the equality in distribution, by ϕ_ξ the characteristic function of the random variable ξ. We will suppose $\sum_{i\in\emptyset} a_i = 0$.

Recall, that $V(t)$, $t \geq 0$ is a Levy process if $V(0) = 0$ and V is a homogeneous stochastically continuous random process with independent increments, such that it's trajectories belong to the Skorohod space. In the paper we will denote by the same symbols the random process and the random element corresponding to this random process.

Using the Levy's representation (see [5], sect. 18) we can assume that the characteristic function of the Levy process $V = V(\gamma, \sigma, L, R)$ is

$$\varphi_{V(t)}(x) = \mathbb{E}\left(e^{ixV(t)}\right) = \psi\left(t, x, L(y), R(y), \sigma, \gamma\right) = \exp\left[itx\gamma - \right.$$

$$\left. -\frac{\sigma^2 x^2 t}{2} + t\left\{\int\limits_{-\infty}^{0}\left(e^{ixy} - 1 - \frac{ixy}{1+y^2}\right)dL(y) + \int\limits_{0}^{\infty}\left(e^{ixy} - 1 - \frac{ixy}{1+y^2}\right)dR(y)\right\}\right],$$

$x \in \mathbb{R}$

Here $L(y)$ is (left-continuous and) non-decreasing on $(-\infty, 0)$ with $L(-\infty) = 0$ and $R(y)$ is (right-continuous and) non-decreasing on $(0, \infty)$ with $R(\infty) = 0$ and

$$\int_{-\varepsilon}^{0} y^2 dL(y) + \int_{0}^{\varepsilon} y^2 dR(y) < +\infty \text{ for all } \varepsilon > 0. \tag{3}$$

Not bounded the community, we can assume $\gamma = 0$ and we will denote $V(\sigma, L, R) = V(0, \sigma, L, R)$.

The main result of the section is the following theorem.

Theorem 1. *Let $V_n = V_n(\sigma_n, L_n, R_n)$, $V = V(\sigma, L, R)$ be Levy processes. Then*

$$V_n \xrightarrow{d} V \text{ as } n \to \infty \tag{4}$$

in $D[0,1]$ if and only if

$$V_n(1) \xrightarrow{d} V(1) \quad \text{as} \quad n \to \infty. \tag{5}$$

Proof. The proof of the necessity is evident. We will prove the sufficiency. By Theorem 2, sect.19 in [5], (5) implies

$$L_n(y) \to L(y), \quad R_n(y) \to R(y), \quad \text{as} \quad n \to \infty$$

and

$$\lim_{\varepsilon\to 0}\overline{\lim_{n\to\infty}}\left\{\int_{-\varepsilon}^{0} u^2 dL_n(u) + \sigma_n^2 + \int_{0}^{\varepsilon} u^2 dR_n(u)\right\} =$$

$$\lim_{\varepsilon\to 0}\underline{\lim_{n\to\infty}}\left\{\int_{-\varepsilon}^{0} u^2 dL_n(u) + \sigma_n^2 + \int_{0}^{\varepsilon} u^2 dR_n(u)\right\} = \sigma^2.$$

Therefore by Theorem 2 of sect. 19 in [5] $V_n(t) \xrightarrow{d} V(t)$ as $n \to \infty$ for all $t \in [0,1]$.

Let $0 \leq t_0 < t_1 < ... < t_n \leq 1$. Introduce the notation $\Delta V_{ni} = V_n(t_i) - V_n(t_{i-1})$ and $\Delta V_i = V(t_i) - V(t_{i-1})$. Since $\Delta V_{ni},\ 1 \leq i \leq k$ are independent random variables and $\Delta V_{ni} \xrightarrow{d} \Delta V_i$ as $n \to \infty$, we have:

$$(\Delta V_{n1}, ..., \Delta V_{nk}) \xrightarrow{d} (\Delta V_1, ..., \Delta V_k) \text{ as } n \to \infty.$$

Consequently, the finite dimensional distributions of V_n converge to the finite dimensional distributions of V.

By Theorem 1 in [11] (p. 229), it's sufficiently to demonstrate that for all $\varepsilon > 0$

$$\lim_{h\to\infty} \overline{\lim_{n\to\infty}} \sup_{|t'-t''|\leq h} P\{|V_n(t') - V_n(t'')| > \varepsilon\}. \tag{6}$$

By Theorem 4 in [12] the set of measures in $D[0,1]$, corresponding to the processes V_n is relatively compact. Then by Theorem 1 in [12] we obtain (6). Proof is completed.

Let $\xi_{ni}, \xi_i\ i \in \mathbf{N}$, be independent identically distributed random variables for all $n \in \mathbb{N}$, $\pi(t)$ is a Poisson random process with the intensity 1, ξ_{ni} and $\pi(t)$ are independent. We will consider the Levy processes

$$V'_n(t) = \sum_{i=1}^{\pi(t)} \xi_{ni},\ t \in [0,1] \tag{7}$$

and

$$V'(t) = \sum_{i=1}^{\pi(t)} \xi_i, \quad t \in [0,1]. \tag{8}$$

Such processes are called the compound processes.

Corollary 1. *Let V'_n and V' be defined by (7), (8). Suppose that*

$$\xi_{ni} \xrightarrow{d} \xi_i \text{ as } n \to \infty. \tag{9}$$

Then we have

$$V'_n \xrightarrow{d} V' \text{ as } n \to \infty \tag{10}$$

in $D[0,1]$.

Proof. Observe that (9) implies

$$\phi_{\xi_{n1}}(x) \to \phi_{\xi_1}(x) \text{ as } n \to \infty \quad \text{for all } x \in \mathbb{R}.$$

Since

$$\phi_{V'_n(1)}(x) = \exp\{\phi_{\xi_{n1}}(x) - 1\}, \quad x \in \mathbb{R}$$

and

$$\phi_{V'(1)}(x) = \exp\{\phi_{\xi}(x) - 1\}, \quad x \in \mathbb{R},$$

we have

$$\phi_{V'_n(1)}(x) \to \phi_{V'(1)}(x) \text{ as } n \to \infty \text{ for all } x \in \mathbb{R}.$$

Consequantly,

$$V'_n(1) \xrightarrow{d} V'(1) \text{ as } n \to \infty$$

and we can apply Theorem 1. Proof is completed.

Consider now the random processes

$$V_n''(t) = \sum_{i=1}^{\pi(k_n t)} \xi_{ni}, \quad t \in [0,1], \quad k_n \in \mathbf{N}, \quad n \in \mathbb{N}. \tag{11}$$

Corollary 2. *Let V_n'' be defined by (11). Suppose that*

$$\sum_{i=1}^{k_n} \xi_{ni} \xrightarrow{d} \gamma \text{ as } n \to \infty, \tag{12}$$

where γ is an infinitely divisible random variable. Then one has

$$V_n'' \xrightarrow{d} V'' \text{ as } n \to \infty \tag{13}$$

in $D[0,1]$, where $V''(t),\ t > 0$ is a Levy process such that $V''(1) \overset{d}{=} \gamma$.

Proof. The convergence (12) implies, that for all $x \in \mathbb{R}$

$$\phi_{\sum_{i=1}^{k_n} \xi_{ni}}(x) \to \phi_\gamma(x) \text{ as } n \to \infty.$$

The characteristic function of $V_n''(1)$ is

$$\phi_{V_n''(1)}(x) = \exp\{k_n(\phi_{\xi_{n1}}(x) - 1)\}, \quad x \in \mathbb{R}.$$

Observe, that

$$k_n \ln \phi_{\xi_{ni}}(x) = k_n \ln(1 - (1 - \phi_{\xi_{ni}}(x))) = (\phi_{\xi_{ni}}(x) - 1)k_n + k_n o(\phi_{\xi_{ni}}(x) - 1).$$

Therefore

$$\phi_{V_n''(1)}(x) \to \phi_{V''(1)}(x) \text{ as } n \to \infty \text{ for all } x \in \mathbb{R}.$$

Consequantly,

$$V_n''(1) \xrightarrow{d} V''(1) \text{ as } n \to \infty$$

and we can apply Theorem 1. Proof is completed.

3 The convergence to α-stable processes.

Let $0 < \alpha < 2$.

Consider the conditions:

$$L(-y)/|R(y)| \to c_1/c_2, \quad \text{as} \quad y \to \infty\,, \tag{14}$$

$$\frac{L(-y) + |R(y)|}{L(-yx) + |R(yx)|} \to x^\alpha, \quad \text{as} \quad y \to \infty\,, \tag{15}$$

where $c_1, c_2 \geq 0,\ c_1 + c_2 > 0$ are valid.

By Theorem 2, sect.35 in [5], the conditions (14), (15) imply that an infinitely divisible random variable with functions $L(y)$ and $R(y)$ in Levy's representation of it's characteristic function belongs to the domain of attraction of the α-stable law having Levy's representation $L_\alpha(y) = \frac{c_1}{|y|^\alpha},\ R_\alpha(y) = \frac{c_2}{-y^\alpha}$. Than it exists $a_n,\ s_n$ such that $s_n,\ a_n \to \infty$ as $n \to \infty$ and

$$\text{for all } y<0 \ \ s_n L(a_n y) \to \frac{c_1}{|y|^\alpha}, \text{ as } n\to\infty, \quad \text{for all } y>0 \ \ s_n R(a_n y) \to \frac{c_2}{-y^\alpha} \text{ as } n\to\infty. \tag{16}$$

Consider the sequence of the random processes

$$X_n(t) = \frac{V(s_n t)}{a_n} - t b_n, \quad t\in[0,1], \quad n\in\mathbb{N}, \tag{17}$$

where $s_n,\ a_n \in \mathbb{R}$ are defined by conditions (16) and

$$b_n = \int_{-\infty}^{0} \frac{z^3(1-a_n^2)}{(1+z^2)(1+a_n^2 z^2)} ds_n L(a_n z) + \int_{0}^{+\infty} \frac{z^3(1-a_n^2)}{(1+z^2)(1+a_n^2 z^2)} ds_n R(a_n z). \tag{18}$$

Denote $Y_\alpha = V(0, \frac{c_1}{|y|^\alpha}, \frac{c_2}{-y^\alpha})$. Then Y_α is a α-stable process $(0<\alpha<2)$. Now we can formulate the next theorem of convergence to a α-stable Levy processes.

Theorem 2. *Let (14) and (15) be valid, s_n and a_n be satisfy (16), X_n be defined by (17), b_n be defined in (18) and $\sigma = 0$. Then we have*

$$X_n \xrightarrow{d} Y_\alpha, \ \ as \ \ n\to\infty$$

in $D[0,1]$.

Proof. The characteristic function of X_n is

$$\varphi_{X_n(t)}(x) =$$

$$= \exp\left[t \left\{ \int_{-\infty}^{0} \left(e^{ixy} - 1 - \frac{ixy}{1+y^2} \right) d(s_n L(a_n y)) + \int_{0}^{\infty} \left(e^{ixy} - 1 - \frac{ixy}{1+y^2} \right) d(s_n R(a_n y)) \right\} \right].$$

Then

$$X_n(1) \xrightarrow{d} Y_\alpha(1)$$

as $n\to\infty$, and by Theorem 1 we obtain the affirmation of theorem. Proof is completed.

Consider the functions

$$L(x) = (c_1 + \alpha_1(x)) \frac{1}{|x|^\alpha}, \quad x<0, \tag{19}$$

$$R(x) = (c_2 + \alpha_2(x)) \frac{1}{-x^\alpha}, \quad x>0, \tag{20}$$

where $\alpha_1(x)$ and $\alpha_2(x)$ are the functions such that

$$\lim_{x\to-\infty} \alpha_1(x) = \lim_{x\to+\infty} \alpha_2(x) = 0. \tag{21}$$

By Theorem 5, sect. 35 in [5], the infinity divisible random variable γ belongs to the domain of normal attraction of the α-stable low if and only if the functions $L(x)$ and $R(x)$ in Levy representation of its characteristic function are (19) and (20), relatively.

Then

$$\text{for all } x<0 \ \ nL(n^{1/\alpha}x) \to \frac{c_1}{|x|^\alpha}, \ \ \text{for all } x>0 \ \ nR(n^{1/\alpha}x) \to \frac{c_2}{-x^\alpha} \ \ \text{as } n\to\infty. \tag{22}$$

Consider the sequence of the random processes

$$X'_n(x) = \frac{V(nt)}{n^{1/\alpha}} - tb'_n, \quad x \in [0,1], n \in \mathbb{N}, \tag{23}$$

where

$$b'_n = \int_{-\infty}^{0} \frac{z^3(1-n^{2/\alpha})}{(1+z^2)(1+n^{2/\alpha}z^2)} dnL(n^{1/\alpha}z) + \int_{0}^{+\infty} \frac{z^3(1-n^{2/\alpha})}{(1+z^2)(1+n^{2/\alpha}z^2)} dnR(n^{1/\alpha}z). \tag{24}$$

Then the next theorem is valid

Theorem 3. *Let X'_n be defined by (23), b_n be defined by (24) and $\sigma = 0$. Then the conditions (19), (20), (21) imply the convergence*

$$X'_n \xrightarrow{d} Y_\alpha, \ \text{as } n \to \infty \tag{25}$$

in $D[0,1]$.

Proof. The characteristic function of X'_n is

$$\varphi_{X'_n(t)}(x) =$$

$$= \exp\left[t \left\{ \int\limits_{-\infty}^{0} \left(e^{ixy} - 1 - \frac{ixy}{1+y^2} \right) dnL(n^{1/\alpha}y) + \int\limits_{0}^{\infty} \left(e^{ixy} - 1 - \frac{ixy}{1+y^2} \right) dnR(n^{1/\alpha}y) \right\} \right].$$

Then

$$X'_n(1) \xrightarrow{d} Y_\alpha(1)$$

as $n \to \infty$, and by Theorem 1 we obtain (25). Proof is completed.

Let now $\alpha = 2$. Consider the condition

$$\frac{x^2 \left(\int_{-\infty}^{-x} dL(y) + \int_x^{+\infty} dR(y) \right)}{\int_{-x}^{0} y^2 dL(y) + \int_0^x y^2 dR(y)} \to 0 \text{ as } x \to \infty. \tag{26}$$

By Theorem 1, sect. 35 in [5], the condition (26) is valid if and only if an infinitely divisible random variable γ with functions $L(y)$ and $R(y)$ in Levy's representation of it characteristic function belongs to the domain of attraction of Gaussian law having Levy's representation $\psi(t, x, 0, 0, \sigma_W)$. Then exists a_n, s_n such that s_n, $a_n \to \infty$ as $n \to \infty$ and

$$\text{for all } y < 0 \ \ s_n L(a_n y) \to 0, \quad \text{for all } y > 0 \ \ s_n R(a_n y) \to 0, \tag{27}$$

$$\lim_{\varepsilon \to 0} \overline{\lim_{n \to \infty}} \left\{ \int_{-\varepsilon}^{0} u^2 ds_n L(a_n u) + \frac{\sigma^2}{a_n^2} s_n + \int_0^{\varepsilon} u^2 ds_n R(a_n u) \right\} =$$

$$\lim_{\varepsilon \to 0} \underline{\lim_{n \to \infty}} \left\{ \int_{-\varepsilon}^{0} u^2 ds_n L(a_n u) + \frac{\sigma^2}{a_n^2} s_n + \int_0^{\varepsilon} u^2 ds_n R(a_n u) \right\} = \sigma_W^2. \tag{28}$$

Consider the sequence of the random processes

$$X_n(t) = \frac{V(s_n t)}{a_n} - tb_n, \quad t \in [0,1], \quad n \in \mathbb{N}, \tag{29}$$

where s_n, $a_n \in \mathbb{R}$ are defined by conditions (27), (28) and

$$b_n = \int_{-\infty}^{0} \frac{z^3(1-a_n^2)}{(1+z^2)(1+a_n^2 z^2)} ds_n L(a_n z) + \int_0^{+\infty} \frac{z^3(1-a_n^2)}{(1+z^2)(1+a_n^2 z^2)} ds_n R(a_n z) -$$

$$- i \lim_{\varepsilon \to 0} \left\{ \int_{-\varepsilon}^{0} z^2 ds_n L(a_n z) + \int_{0}^{\varepsilon} z^2 ds_n R(a_n z) \right\}. \tag{30}$$

Theorem 4. *Let a_n, s_n be satisfied (27),(28) and X_n be defined by (29), b_n be defined by (30). Then the condition (26) implies*

$$X_n \xrightarrow{d} \sigma_W W, \ \ as \ \ n \to \infty,$$

in $D[0,1]$.

Proof. The characteristic function of $X_n(t)$ is:

$$\varphi_{X_n(t)}(x) = \exp \left[-s_n t \frac{\sigma^2}{2a_n^2} x^2 + \right.$$

$$\left. +t \left\{ \int_{-\infty}^{0} (e^{ixy} - 1 - \frac{ixy}{1+y^2}) ds_n L(a_n y) + \int_{0}^{\infty} (e^{ixy} - 1 - \frac{ixy}{1+y^2}) ds_n R(a_n y) \right\} \right].$$

Then

$$X_n(1) \xrightarrow{d} \sigma_W W(1)$$

as $n \to \infty$. Then by Theorem 1

$$X_n \xrightarrow{d} \sigma_W W.$$

Proof is completed.

Consider the random processes

$$X_t(x) = \frac{V(tx)}{f(t)} - x g(t), \ \ x \in [0,1], t \in (0, \infty), \tag{31}$$

$$g(t) = \int_{-\infty}^{0} \frac{v^3 (f(t)^2 - 1)}{f(t)(1+v^2)(f(t)^2 + v^2)} dt L(f(t)v) + \int_{0}^{\infty} \frac{v^3 (f(t)^2 - 1)}{f(t)(1+v^2)(f(t)^2 + v^2)} dt R(f(t)v) \tag{32}$$

4 Almost sure versions.

Now we will consider the sequence of measures $Q_n(\omega)$, defined in (2) and connected with the random processes X_n, defined in (1) with the condition for sequence s_n:

(A) for some $\beta > 0$ the sequence $\left(\frac{s_n}{n^\beta} \right)$ is increasing as $n \to \infty$.

Let Y_α is a α-stable random process ($0 < \alpha \leq 2$). We will denote by $\xrightarrow{w}$ the weak convergence of measures, by μ_α the distribution of the random element Y_α and by $\rho(x,y)$ the usual metric in $D[0,1]$, with which $D[0,1]$ is a separable complete space.

Let ξ_{ni} are independent identically distributed random variables. Consider the sums $S_n = \sum_{i=1}^{k_n} \xi_{ni}$. Below we will used the next lemma:

Lemma 1. *Let S be an infinitely divisible random variable such that*

$$S_n \xrightarrow{d} S \ \ as \ n \to \infty. \tag{33}$$

Then for all $C > 0$ the next conditions are valid:

$$C_1 = \sup_{n\in\mathbf{N}} k_n P\{|\xi_{n1}| \geq C\} < +\infty,$$

$$C_2 = \sup_{n\in\mathbf{N}} k_n E(\xi_{n1} I_{\{|\xi_1|<C\}}) < +\infty,$$

$$C_3 = \sup_{n\in\mathbf{N}} k_n D(\xi_{n1} I_{\{|\xi_{n1}|<C\}}) < +\infty,$$

Proof. By Theorem 1, sect. 23 in [5], the convergence (33) implies that

$$\sup_{n\in\mathbf{N}} k_n E \frac{\xi_{n1}^2}{1+\xi_{n1}^2} < +\infty.$$

Observe, that

$$\sup_{n\in\mathbf{N}} k_n P\{|\xi_{n1}| \geq C\} = \sup_{n\in\mathbf{N}} k_n E I_{\{|\xi_{n1}|\geq C\}} = \sup_{n\in\mathbf{N}} k_n \frac{C^2+1}{C^2} E\left(\frac{C^2}{1+C^2} I_{\{|\xi_{n1}|\geq C\}}\right) \leq$$

$$\leq \sup_{n\in\mathbf{N}} k_n \frac{C^2+1}{C^2} E\left(\frac{\xi_{n1}^2}{1+\xi_{n1}^2}\right) < +\infty;$$

By Theorem 2 in [13] (p. 373),

$$\sup_{n\in\mathbf{N}} k_n E(\xi_{n1} I_{\{|\xi_{n1}|<C\}}) < +\infty;$$

$$\sup_{n\in\mathbf{N}} k_n D\left(\xi_{n1} I_{\{|\xi_{n1}|<C\}}\right) \leq \sup_{n\in\mathbf{N}} k_n E\xi_{n1}^2 I_{\{|\xi_{n1}|<C\}} =$$

$$= \sup_{n\in\mathbf{N}} k_n (1+C^2) E\left(\frac{\xi_{n1}^2}{1+C^2} I_{\{|\xi_{n1}|<C\}}\right) \leq \sup_{n\in\mathbf{N}} k_n (1+C^2) E\left(\frac{\xi_{n1}^2}{1+\xi_{n1}^2}\right) < +\infty.$$

The proof is completed.

Theorem 5. *Let (A) be valid and $X_n \xrightarrow{d} Y_\alpha$ as $n \to \infty$. Then it holds*

$$Q_n(\omega) \xrightarrow{w} \mu_\alpha \tag{34}$$

as $n \to \infty$ for almost all $\omega \in \Omega$ in $D[0,1]$.

Proof. Let $0 < l < k$ and

$$X_{lk}(t) = \begin{cases} b_k t & 0 \leq t \leq \frac{s_l}{s_k} \\ X_k(t) - \frac{V(s_l)}{a_k} & \frac{s_l}{s_k} \leq t \leq 1 \end{cases}$$

Then X_{lk} and X_l are independent random processes.

It's clear, that $D[0,1]$ is a complete separable space in the metric

$$\rho_0(x,y) = \frac{\rho(x,y)}{1+\rho(x,y)}, \quad x, y \in D[0,1].$$

We will consider the metric

$$\rho_1(x,y) = \min\{\sup_{0\leq t\leq 1} |x(t)-y(t)|, 1\}.$$

Observe, that

$$\rho_1(x,y) \geq \rho_0(x,y), \quad x,\ y\ \in D[0,1].$$

We will show that

$$E\rho_1(X_k, X_{kl}) \leq C\left(\frac{l}{k}\right)^{\frac{\beta}{2}}. \tag{35}$$

Since $V(t)$ is a random process with independent increments,

$$V(s_l) = \sum_{i=1}^{n_l} \xi_i,$$

where $\xi_i = V\left(\frac{s_l i}{n_l}\right) - V\left(\frac{s_l(i-1)}{n_l}\right)$ are independent identically distributed random variables. Let $n_k \in \mathbf{N}$ such that $\frac{n_l}{n_k} \leq \frac{s_l}{s_k}$ and it exists $\lambda \in [0,1]$ such that

$$V(\lambda) = \sum_{i=1}^{n_k} \xi_i.$$

We have

$$E\min\left\{\frac{1}{a_k}\sum_{i=1}^{n_l}|\xi_i I_{\{|\xi_i| \geq a_k\}}|, 1\right\} \leq n_l E\min\{\frac{1}{a_k}|\xi_i| I_{\{|\xi_i| \geq a_k\}}, 1\} \leq n_l P\{|\xi_1| \geq a_k\} =$$

$$= \frac{n_l}{n_k} n_k P\{|\xi_1| \geq a_k\} \leq C_1\left(\frac{l}{k}\right)^{\beta} \leq C_1\left(\frac{l}{k}\right)^{\frac{\beta}{2}},$$

and

$$E\min\left\{\frac{1}{a_k}\left|\sum_{i=1}^{n_l}\xi_i I_{\{|\xi_i|<a_k\}}\right|, 1\right\} =$$

$$E\min\{\frac{1}{a_k}|\sum_{i=1}^{n_l}\xi_i I_{\{|\xi_i|<a_k\}} - E\xi_i I_{\{|\xi_i|<a_k\}} + E\xi_i I_{\{|\xi_i|<a_k\}}|, 1\} \leq$$

$$\frac{s_l}{s_k}\sqrt{n_k D(\frac{1}{a_k}\xi_1 I_{\{|\xi_1|<a_k\}})} + \frac{s_l}{s_k} n_k E(\frac{1}{a_k}\xi_1 I_{\{|\xi_1|<a_k\}}) \leq (C_2 + \sqrt{C_3})\left(\frac{l}{k}\right)^{\frac{\beta}{2}},$$

where $C_1,\ C_2,\ C_3$ are defined in Lemma 1.

Then, using the moment inequality from [8], sect. 5, p. 231 we obtain

$$E\rho_1(X_k, X_{lk}) = E\min\{\sup_{0\leq t\leq 1}|X_k - X_{lk}|, 1\} = E\min\left\{\sup_{0\leq t\leq \frac{s_l}{s_k}}|\frac{V(ts_k)}{a_k}|, 1\right\} \leq$$

$$4E\min\left\{\left|\frac{V(s_l)}{a_k}\right|, 1\right\} = 4E\min\{\frac{1}{a_k}|\sum_{i=1}^{n_l}\xi_i|, 1\} \leq$$

$$4E\min\{\frac{1}{a_k}|\sum_{i=1}^{n_l}\xi_i I_{\{|\xi_i|<a_k\}}|, 1\} + 4E\min\{\frac{1}{a_k}|\sum_{i=1}^{n_l}\xi_i I_{\{|\xi_i|\geq a_k\}}|, 1\} \leq 4(C_1 + C_2 + \sqrt{C_3})\left(\frac{l}{k}\right)^{\frac{\beta}{2}}.$$

Then by Lemma 1 from [2], this and Theorems 2-4 imply the convergence (34). The proof is completed.

References.

[1] M. T. Lacey and W. Philipp, A note on almost sure central limit theorem, *Statistics and Probability Letters* **9**(2) (1990), 201–205.

[2] A. Chuprunov and I. Fazekas, Almost sure versions of some analogues of the invariance principle, *Publicationes Mathematicae, Debrecen* **54**(3-4) (1999), 457–471.

[3] A. Chuprunov and I. Fazekas, Almost sure limit theorems for the Pearson statistic, *Teor. Veroyatnost. i Primenen.* **48**(1), 162–169.

[4] A. Chuprunov and I. Fazekas, Integral analogues of almost sure limit theorems, *Periodica Mathematica Hungaricao* V.5 (1-2), 2005 pp. 61-78.

[5] B. V. Gnedenko and A. N. Kolmogorov, Limit Distributions for Sums of Independent Random Variables, *Addison-Wesley, Reading, Massachusetts*, 1954.

[6] P. Billingsley, Convergence of probability measures, *John Wiley and Sons, New York*, 1968.

[7] D. Pollard, Convergence of stochastic processes, *Springer-Verlag, New York*, 1984.

[8] N. N. Vakhania, V. I. Tarieladze and S. A. Chobanian, Probability distributions in Banach spaces, *Nauka, Moscow*, 1985 (in Russian).

[9] V. M. Kruglov and V. Yu. Korolev, Limit theorems for random sums, *Moscow University Press, Moscow*, 1990 (in Russian).

[10] I. I. Gihman, A. V Skorohod, Theory of the random processes, *Nauka, Moscow*, 1973 (in Russian).

[11] A. V. Skorohod, Random processes with independent increments, *Probability theory and mathematical statistics, Nauka, Moscow*, 1964, (in Russian).

[12] V. M. Borodikhin, Conditions for the compactness of subsets of measures on some metric spaces, *Teor. Veroyat. i Matem. Statist.* **3** (1970), 16-28, (in Russian).

[13] A. N. Shiryaev, Probability, *Nauka, Moscow*, 1997 (in Russian)